

those from neighboring islands. The wings and tail are very dark, almost black, but my cabinet also contains specimens from Porto Rico which are difficult to distinguish from birds taken in Cuba and San Domingo.

Mimus polyglottos portoricensis BRYANT.

Mimus polyglottos var. *portoricensis* BRYANT. Proc. Boston Soc. Nat. Hist. XI, p. 68 (1866).

Mimus polyglottos GUNDLACH, Anales de la Soc. Esp. de Hist. Nat. VII, p. 173 (1878).

HABITAT.—Porto Rico.

A COLLECTION OF BIRDS TAKEN BY CYRUS S. WINCH IN THE ISLANDS OF ANGUILLA, ANTIGUA, AND ST. EUSTATIUS, WEST INDIES, DURING APRIL, MAY, JUNE, AND A PART OF JULY, 1890.

BY CHARLES B. CORY.

ANGUILLA.

<i>Podilymbus podiceps</i> (<i>Linn.</i>). <i>Gallinula galeata</i> (<i>Licht.</i>). <i>Fulica caribaea</i> (<i>Ridgw.</i>). <i>Himantopus mexicanus</i> <i>Müll.</i> <i>Falco caribbearum</i> <i>Gmel.</i> <i>Ceryle alcyon</i> (<i>Linn.</i>). <i>Tyrannus dominicensis</i> (<i>Gmel.</i>).	<i>Elænea martinicana</i> (<i>Linn.</i>). <i>Euetheia bicolor</i> (<i>Linn.</i>). <i>Loxigilla noctis</i> (<i>Linn.</i>). <i>Cæreba bartholemica?</i> (<i>Sparrm.</i>). <i>Cichlherminia fuscata densirostris</i> (<i>Vieill.</i>). <i>Galeoscoptes carolinensis</i> (<i>Linn.</i>).
---	---

ANTIGUA.

<i>Larus atricilla</i> (<i>Linn.</i>). <i>Sterna sandvicensis acuflavida</i> <i>Cabot.</i> <i>Sterna dougalli</i> <i>Mont.*</i> <i>Sterna antillarum</i> (<i>Less.</i>).	<i>Pelecanus fuscus</i> <i>Linn.</i> <i>Gallinula galeata</i> (<i>Licht.</i>). <i>Fulica caribaea</i> <i>Ridgw.</i> <i>Himantopus Müll.</i>
---	--

* Specimens of this species taken in June have the basal half of the bill red, and in three specimens the red color extends more than half the length of the bill. In all these birds the black cap is complete. In specimens taken in September, which have the black cap imperfect or mixed with white and gray, the bill is entirely black.

Symphearia semipalmata (<i>Gmel.</i>). Totanus melanoleucus (<i>Gmel.</i>). Colinus virginianus (<i>Linn.</i>). Columba leucocephala <i>Linn.</i> . Zenaida zenaida (<i>Bonap.</i>). Geotrygon mystacea (<i>Temm.</i>). Columbigallina passerina (<i>Linn.</i>). Buteo latissimus (<i>Wils.</i>). Falco caribbearum (<i>Gmel.</i>). Speotyto amanra <i>Lafr.</i> Coccyzus minor (<i>Gmel.</i>). Coccyzus americanus (<i>Linn.</i>).	Bellona exilis (<i>Gmel.</i>) Eulampis holosericeus (<i>Linn.</i>). Tyrannus dominicensis (<i>Gmel.</i>). Elæna martinica (<i>Linn.</i>). Euetheia bicolor (<i>Linn.</i>). Loxigilla noctis (<i>Linn.</i>). Piranga erythromelas <i>Vieill.</i> Vireo calidris (<i>Linn.</i>). Cæreba dominicana (<i>Taylor</i>). Dendroica ruficapilla (<i>Gmel.</i>). Allenia montana (<i>Lafr.</i>). Cichlherminia fuscata densirostris (<i>Vieill.</i>).
--	--

ST. EUSTATIUS.

Phaëthon flavirostris <i>Brandt</i> . Ardea cœrulea <i>Linn.</i> . Columba corensis <i>Gmel.</i> . Zenaida zenaida (<i>Bonap.</i>). Falco caribbearum <i>Gmel.</i> . Bellona exilis (<i>Gmel.</i>). Eulampis holosericeus (<i>Linn.</i>). Tyrannus dominicensis (<i>Gmel.</i>). Euetheia bicolor (<i>Linn.</i>). Loxigilla noctis (<i>Linn.</i>).	Vireo calidris (<i>Linn.</i>). Progne dominicensis (<i>Gmel.</i>). Cæreba dominicana (<i>Taylor</i>). Mniotila varia (<i>Linn.</i>). Dendroica ruficapilla (<i>Gmel.</i>). Compsothlypis americana (<i>Linn.</i>). Dendroica discolor (<i>Vieill.</i>). Setophaga ruticilla (<i>Linn.</i>). Allenia montana (<i>Lafr.</i>). Cichlherminia fuscata densirostris (<i>Vieill.</i>).
--	--

A LIST OF THE BIRDS COLLECTED IN THE
ISLANDS OF ST. CROIX AND ST. KITTS, WEST
INDIES, DURING MARCH AND APRIL, AND
IN GUADELOUPE DURING AUGUST,
SEPTEMBER, AND OCTOBER, 1890.

BY CHARLES B. CORY.

ST. CROIX.

Gallinula galeata (<i>Licht.</i>). Ardea virescens <i>Linn.</i> . Nycticorax violaceus (<i>Linn.</i>).	Ctanus flavipes (<i>Gmel.</i>). Columba leucocephala <i>Linn.</i> . Columba corensis <i>Gmel.</i>
--	---

* A rather small pale form, resident in Antigua, and quite abundant.